



BIOSpain
2012

6th International Meeting on Biotechnology
19 - 21 September, 2012 - Bilbao (Spain)

SCIENTIFIC CONGRESS
BIOTEC 2012



ORGANIZED BY:



SPONSOR:



CO-SPONSOR:



S-8. OC1. Omics and Systems Biotechnology

Profiling and structural analysis of *Ashbya gossypii* N-glycansT. Q. Aguiar¹, H. Maaheimo², A. Heiskanen³, M. Wiebe², M. Penttilä² and L. Domingues¹

¹IBB - Institute for Biotechnology and Bioengineering, Centre of Biological Engineering, Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal, luciliad@deb.uminho.pt

²VTT Technical Research Centre of Finland, P.O. Box 1000, 02044 VTT Espoo, Finland

³Glykos Finland Ltd., Helsinki, Finland

Ashbya gossypii is a filamentous hemiascomycete used for the industrial production of riboflavin (vitamin B2) that has recently been explored as a new fungal host system for the expression and secretion of proteins. However, the secretory pathway of *A. gossypii* remains poorly characterized, although evidences indicate that the level of secretion in *A. gossypii* is more similar to yeast than to other filamentous fungi; and that recombinant glycoproteins produced by *A. gossypii* are less extensively glycosylated than those produced by *Saccharomyces cerevisiae*. A better understanding of the *A. gossypii* glycobiology offers a basis to characterize its protein secretory processes and may possibly be of practical impact in the production of heterologous proteins.

As a starting point on the elucidation of *A. gossypii* N-glycosylation, in this study we performed a global analysis of whole mixtures of N-glycans attached to *A. gossypii* native secreted glycoproteins. Through combined analysis of N-glycan spectra obtained by MALDI-TOF mass spectrometry and ¹H NMR spectroscopy we report here that the major N-glycan species derived from *A. gossypii* secreted proteins are high-mannose type glycans containing core-type structures (Man8-9GlcNac2). Smaller core structures (Man5-7GlcNac2) are also present in lower abundance and N-glycans with long outer chains are only present in minor relative amounts. All identified acidic N-glycans contain a phosphate residue. Among the most abundant

acidic structures are short phosphorylated outer chain glycans (PhMan13-15GlcNac2).

Our data present the first report on the composition, size distribution and structural characteristics of *A. gossypii* N-glycans.

Acknowledgments: This work was supported by Fundação para a Ciência e a Tecnologia (FCT), Portugal: Project AshByofactory grant PTDC/EBB-EBI/101985/2008 and MIT-Portugal Program PhD grant SFRH/BD/39112/2007 to Tatiana Quinta Aguiar. The authors would like to thank Olli Aitio for helping with the interpretation of the NMR spectra.